WHAT IS CLAIMED IS:

A C-11 protein comprising a protein selected fom the group consisting of:

- (a) a protein comprising/an amino acid sequence having SEQ ID NO. 2 (SEQUENCE LISTING); and
- (b) a protein comprising an amino acid sequence that is derived from the amino acid sequence having SEQ ID NO. 2 by deletion, substitution or insertion of one or more amino acids / said protein having cellcalcification inhibitory activity.
- 2. A gene encoding the protein according to claim 1.
- A pharmaceutical composition comprising the protein according to, claim 1.
- The pharmaceutical composition according to claim 3, wherein said composition is intended for a cell-calcification inhibitor.
 - A cell-calcification inhibitor comprising a cg protein selected from the group consisting of:
- (a) a protein comprising an amino acid sequence having SEQ ID NO. 4 (SEQUENCE/LISTING); and
- (b) a protein comprising an amino acid sequence that is derived from the amino acid sequence having SEQ ID NO. 4 by deletion, substitution or insertion of one or more amino acids, said protein having cellcalcification inhibitory activity.

10

5

15

20

5

- 6. An antibody to the C-11 protein according to claim 1.
- 7. The antibody according to claim 6, wherein said antibody is a monoclonal antibody.
- 8. A method for measuring the calcification of cells comprising:

measuring the expression of a C-11 gene or a c-erg gene in the cells.

- 9. The method according to claim 8, wherein the expression of the gene is measured by the amount of C-11 mRNA expressed in the cells or the amount of c-erg mRNA expressed in the cells using a probe against a DNA sequence specific to the C-11 gene or to the c-erg gene.
- 10. The method according to claim 8, wherein the expression of the gene is measured by the amount of expression of a C-11 protein in the cells or the amount of expression of a c-erg protein in the cells.
- expression of the gene is measured by the amount of the C-11 protein expressed in the cells or the amount of the c-erg protein expressed in the cells by means of the antibody according to claims 6, or 7.
- 12. A method for diagnosing osteoarthritis or OPLL comprising:

measuring the cell-calcification using a method

25

15

5

according to any of claims 8-11.

- 13. A kit for measuring the cell-calcification of cells comprising either or both of an antibody to a C-11 protein and an antibody to a c-erg protein.
- 14. A method for screening a substance having cell-calcification inhibitory blocking activity, said method comprising using cells transformed with a gene encoding a protein selected from the group consisting of:
- (a) a protein comprising an amino acid sequence having SEQ ID NO. 2 (SEQUENCE LISTING);
- (b) a protein comprising an amino acid sequence that is derived from the amino acid sequence having SEQ ID NO. 2 by deletion, substitution or insertion of one or more amino acids, said protein having cell-calcification inhibitory activity;
- (c) a protein comprising an amino acid sequence having SEQ ID NO. 4. (SEQUENCE LISTING); and
- (d) a protein comprising an amino acid sequence that is derived from the amino acid sequence having SEQ ID NO. 4 by deletion, substitution or insertion of one or more amino acids, said protein having cell-calcification inhibitory activity.
- 15. A pharmaceutical composition comprising an erg protein.
 - 16. A pharmaceutical composition comprising an

20

erg gene.

- 17. A pharmaceutical composition comprising a C-11 protein or a c-erg protein.
- A pharmadeutical composition comprising a C-11 gene or a cterg gene.
- A pharmaceutical composition comprising a protein having a consensus amino acid sequence between a c-erg protein and a C-11 protein.